

Obligatory tests inscribed in the French Stud Book for the 2009 stud season.

Tests for CEM and EVA to be taken after the 1st of January and before covering

	Mares before covering			Stallion
	MAIDEN	Barren	In Foal / with foal at foot	Before requesting covering authorisation from H.N.
EQUINE CONTAGIOUS METRITIS	Area : clitoris			Area : urethral fossa
	1 negative result : ok	1 negative result: ok	1 negative result: ok	1 negative result : ok 1 negative result 15 days after the last service and before October 1st: ok
EQUINE VIRAL ARTERITIS	1 serological negative result : ok 2 serological stable or declining positive results at a minimum of 14 day interval : ok 1 serological stable or declining positive result at the start of the year N in compa (serologies in the same laboratory)			1 serological negative result : ok 1 serological positive result with correct vaccination certificate: ok 1 serological positive result without the correct vaccination certificate : - virology in the sperm twice at a 7 day interval with a negative result : ok laboratory) - positive virology, shedder stallion =>specific protocol
EQUINE INFECTIOUS ANEMIA				Coggins test : 1 not more than 3 months old with negative result in order to be admitted to pu covering, and then a test every 3 years

Obligatory vaccination in the sanitary annex rules of the 2009 French Stud Book

INFLUENZA	Before covering	Before requesting covering authorisation from H.N.
	Either a primo vaccination : 2 injections separated by 3 weeks to 3 months plus an annual boost Either an annual booster over the three previous years (with an 8 day tolerance)	Either a primo vaccination : 2 injections separated by 3 weeks to 3 months, plus an annual boost Either an annual booster over the three previous years (with an 8 day tolerance)
Any rupture of the vaccination protocol necessitates a primo vaccination		
Rhinopneumonitis EHV1.4	Before covering	Before requesting the covering authorisation from H.N.
	last injection less than a year primovaccination : 2 injections separated by 3 to 6 weeks At least 1 annual booster, best if 2 a year Or at least 1 annual injection over the last 3 years	last vaccination less than 1 year before obtaining the official covering authorisation. primovaccination : 2 injections separated by 3 to 6 weeks At least 1 annual booster, best if 2 a year Or at least 1 annual injection over the last 3 years
Any rupture of the vaccination protocol necessitates a primo vaccination		

Facultative vaccination

EQUINE VIRALE ARTERITIS		ARTERVAC® Negative serology and then a primo vaccination of 2 injections separated by 3 to 6 weeks on the stallion maintained in isolation Booster every 6 months (with a tolerance of 2 months)
		<i>In the instance of vaccination breakdown :</i> Negative serology and then a primo vaccination of 2 injections separated by 3 to 6 weeks, booster every 6 months (with a tolerance of 2 months)
		Positive serology : virology in the sperm 1 (for 2007) with a negative result, vaccination boost Booster every 6 months (with a tolerance of 2 months)

Cette fiche peut être consultée et téléchargée sur le site www.syndicatdeseleveurs.org